

D-CYCLOSERINE ANTIMICROBIC SUPPLEMENT

VIAL CONTENTS (for 500 ml of medium)

D-Cycloserine.....200 mg

DIRECTIONS

TSC Agar (ISO formulation): aseptically reconstitute the contents of one vial of D-Cycloserine Antimicrobial Supplement with 5 ml of sterile distilled water. Prepare 500 ml of TSC Agar Base (402158), cooled to 45-50°C. Add the supplement, mix well and distribute into sterile Petri dishes.

TSC Agar: aseptically reconstitute the contents of one vial of D-Cycloserine Antimicrobial Supplement with 5 ml of sterile distilled water. Prepare 500 ml of Clostridium Perfringens Agar Base (401307), cooled to 45-50°C. If required by laboratory procedures add 50 ml/litre of Egg Yolk Emulsion 50% (42111601). Add the supplement, mix well and distribute into sterile Petri dishes

INTENDED USE

For the selective isolation and enumeration of *Clostridium perfringens* in foods. For the details of the procedures see the technical sheets of TSC Agar Base, cat. n° 402158 and Clostridium Perfringens Agar Base, cat. N° 401307.

STORAGE

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

PRECAUTIONS AND WARNING

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

PACKAGING

4240002 D-Cycloserine Antimicrobial Supplement 10 vials, each for 500 ml of medium