

## LISTERIA OXFORD ANTIMICROBIC SUPPLEMENT

### VIAL CONTENTS (for 500 ml of medium)

Cicloheximide	200.0 mg
Colistin sulphate	10.0 mg
Cefotetan	1.0 mg
Fosfomicine	5.0 mg
Acriflavine	2.5 mg

### DIRECTIONS

Dissolve the contents of one vial with 5 ml of sterile distilled water under aseptic conditions.

Add to 500 ml of Listeria Oxford Agar Base (cat. N° 401600), autoclaved and cooled to 50°C, Mix well and pour into sterile Petri dishes.

### INTENDED USE

The selective supplement here described is used for the preparation of OXFORD plating medium for the isolation of *Listeria* spp. from food, environmental or clinical specimens. For the details of the procedure, see the technical sheet of Listeria Oxford Agar Base (cat. N° 401600).

### STORAGE

Store at 2-8° - When stored as directed the supplement remains stable until the expiry date shown on the label. Do not use beyond stated expiry date.

### PRECAUTIONS AND WARNING

For *in vitro* diagnostic use only. The supplement should be used only by adequately trained personnel with knowledge of microbiological techniques in the laboratory.

Consult the material safety data sheet before the use.

### PACKAGING

**4240038 Listeria Oxford Antimicrobial Supplement, 10 vials, each for 500 ml of medium**